

TBEV ImmuneSelect Peptide Pool Premium

Catalogue	1292-07
Components	7 nmol/peptide (approx. 11.6 µg), for the stimulation of 1×10^8 cells.
Purity grade	≥90% (HPLC)
Format	Lyophilized, no additive
Storage	Once reconstituted in sterile water, store aliquots at -20 °C or below. Aliquots are stable for 6 months. Do not add salts or buffers to the storage solution, as this may affect peptide stability.

Description

Tick-borne encephalitis virus (TBEV) ImmuneSelect peptide pool is composed of 46 peptides derived from

- ◆ Genome polyprotein (UniProt: P14336).

The peptides include both MHC-II epitopes and the purity of each peptide is ≥90% (HPLC-MS).

Application

The *in vitro* stimulation of Peripheral Blood Mononuclear Cells (PBMCs) with TBEV ImmuneSelect Peptide Pool activates memory T cells specific to the virus, leading to cytokine secretion and the increased expression of specific surface markers.

Reconstitution of peptide pool

1. Allow the lyophilized peptide pool vial to equilibrate to room temperature before opening.
2. Add 233 µL of sterile, nuclease-free water directly to the vial to fully dissolve the contents.
3. Vortex thoroughly to ensure complete solubilization of peptides.
4. The resulting stock solution contains each peptide at a concentration of 30 nmol/mL.
5. Prepare single-use working aliquots to avoid repeated freeze-thaw cycles.
6. Store working aliquots at -20°C.

Recommendations for *in vitro* stimulation

1. Wash PBMCs with sterile cell culture medium, centrifuge at $300 \times g$ for 10 min, and aspirate the supernatant.
2. Resuspend PBMCs in appropriate cell culture medium at 4×10^6 – 1×10^7 cells/mL depending on the downstream experiment. For ELISpot assays, we recommend plating 200,000 cells/well in a 96-well plate.
3. Mix the reconstituted peptide stock thoroughly before use. Dilute the peptide stock solution in sterile cell culture medium to $2 \times$ the final working concentration (1.2 nmol/mL) and sterile-filter the solution prior to use.
4. Add the peptide solution to the wells at a 1:1 (v/v) ratio with the cell suspension, resulting in a final peptide concentration of 0.6 nmol/mL per peptide.
5. Incubate cells for desired time depending on the application.

Note: Incubation times can differ depending on the application. For IFN-γ ELISpot assays, an incubation period of 18–48 hours is recommended.

Need assistance?

Contact our technical support team at support@viraxbiolabs.com
We are happy to help via email or can schedule a call on request